

DYNAMICAL MORDELL-LANG CONJECTURE FOR BIRATIONAL POLYNOMIAL MORPHISMS ON $\mathbb{A}^2$

XIE JUNYI

ABSTRACT. We prove the dynamical Mordell-Lang conjecture for birational polynomial morphisms on $\mathbb{A}^2$.

1. INTRODUCTION

The Mordell-Lang conjecture proved by Faltings [10] and Vojta [21] says that if V is a subvariety of a semiabelian variety G defined over $\mathbb{C}$ and Γ is a finitely generated subgroup of $G(\mathbb{C})$, then $V(\mathbb{C}) \cap \Gamma$ is a union of at most finitely many translates of subgroup of Γ .

The following dynamical analogue of the Mordell-Lang conjecture was proposed by Ghioca and Tucker.

Dynamical Mordell-Lang Conjecture ([14]). Let X be a quasiprojective variety defined over $\mathbb{C}$, let $f : X \rightarrow X$ be an endomorphism, and V be any subvariety of X . For any point $p \in X(\mathbb{C})$ the set $\{n \in \mathbb{N} \mid f^n(p) \in V(\mathbb{C})\}$ is a union of at most finitely many arithmetic progressions.

An arithmetic progression is a set of the form $\{an + b \mid n \in \mathbb{N}\}$ with $a, b \in \mathbb{N}$ possibly with $a = 0$.

Observe that this conjecture implies the classical Mordell-Lang conjecture in the case $\Gamma \simeq (\mathbb{Z}, +)$.

The Dynamical Mordell-Lang conjecture has been proved by Denis [7] for automorphisms of projective spaces and was later generalized by Bell [4] to the case of automorphisms of affine varieties. In [3], Bell, Ghioca and Tucker proved it for étale maps of quasiprojective varieties. The conjecture is also known in the case where $f = (F(x_1), G(x_2)) : \mathbb{A}_{\mathbb{C}}^2 \rightarrow \mathbb{A}_{\mathbb{C}}^2$ and in the case where F, G are polynomials and the subvariety V is a line ([15]), and in the case where $f = (F(x_1), \dots, F(x_n)) : \mathbb{A}_K^n \rightarrow \mathbb{A}_K^n$ if $F \in K[t]$ (for a number field K) is an indecomposable polynomial which has no periodic critical points other than the point at infinity ([5]) and V is a curve.

Our aim is to prove the Dynamical Mordell-Lang conjecture for birational polynomial morphisms on $\mathbb{A}^2$.

Theorem A. *Let K be any algebraically closed field of characteristic 0, and $f : \mathbb{A}_K^2 \rightarrow \mathbb{A}_K^2$ be any birational polynomial morphism defined over K . Let C be*

any curve in $\mathbb{A}_K^2$, and p be any point in $\mathbb{A}^2(K)$. Then the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is a union of at most finitely many arithmetic progressions.

In the case the map is an automorphism of $\mathbb{A}_K^2$ of Hénon type (see [12]) the this result follows from [3]. Our proof provides however an alternative approach and does not rely on the construction of p -adic invariant curves.

Recall that the *algebraic degree* of a polynomial transformation $f(x, y) = (f_1(x, y), f_2(x, y))$ is defined by $\deg f := \max\{\deg f_1, \deg f_2\}$. The limit $\lambda(f) := \lim_{n \rightarrow \infty} (\deg f^n)^{1/n}$ exists and we refer to it as the *dynamical degree* of f (see [8, 9]). Our proof shows that when $\lambda(f) > 1$, then Theorem A holds for fields of arbitrary characteristic.

Note however that our Theorem A does not hold when $\text{char} K > 0$ and $\lambda(f) = 1$ (see [4, Proposition 6.1] for a counter-example).

Let us explain our strategy to prove Theorem A. By some reduction arguments, we may assume that $K = \overline{\mathbb{Q}}$.

If we exclude some special cases, for a birational polynomial morphism $f : \mathbb{A}^2 \rightarrow \mathbb{A}^2$, we may compactify $\mathbb{A}^2$ to a smooth projective surface, such that f extends to a birational transformation on X and there is a fixed point Q in X such that f contracts all curves at infinity to Q (see [11] and Section 6.1).

Our idea is to take advantage of this attracting fixed point and to apply the following local version of the Dynamical Mordell-Lang conjecture.

Theorem 1.1. *Let X be a projective surface over an arbitrary valued field $(K, |\cdot|)$ and $f : X \dashrightarrow X$ be a birational transformation defined over K . Let C be any curve in X and p be any point such that $f^n(p) \in X \setminus I(f)$ for all integers $n \geq 0$ and $f^n(p)$ tends to a fixed point $Q \in I(f^{-1}) \setminus I(f)$ with respect to a projective metric induced by $|\cdot|$ on X .*

If the set

$$\{n \in \mathbb{N} \mid f^n(p) \in C\}$$

is infinite, then either $f^n(p) = Q$ for some $n \geq 0$ or C is fixed.

We now rely on a global argument. When the curve C is passing through the fixed point Q in X , we cover the $\overline{\mathbb{Q}}$ -points of the curve C by the basin of attraction of Q with respect to all absolute values on $\overline{\mathbb{Q}}$. If the point p belongs to one of these attracting basins, then the local dynamical Mordell-Lang Theorem applies and we are done. Otherwise it is possible to bound the height of p and Northcott theorem shows that it is periodic.

When neither the curve C nor its iterates contain the fixed point Q , then we are in position to apply the next result which allows us to conclude.

Theorem 1.2. *Let X be a smooth projective surface over an algebraically closed field, $f : X \dashrightarrow X$ be an algebraically stable birational transformation and C be an irreducible curve in X such that f^n does not contract C for any n . If $f^n(C) \cap I(f) \neq \emptyset$ for all n , then C is periodic.*

The article is organized in 8 sections. In Section 2 we give background informations on surface birational maps and metrics on projective varieties defined

over a valued field. In Section 3 we prove the Theorem 1.2, which is a criterion for a curve to be periodic. In Section 4 we define the *DML* property and prove some basic properties of it. In Section 5 we prove Theorem 1.1 the local version of the Dynamical Mordell-Lang conjecture. In Section 6 we prove Theorem A in the case the dynamical degree $\lambda(f) = 1$. In Section 7 we prove a technical lemma which gives a up bound on height when $\lambda(f) > 1$. In Section 8 we prove Theorem A.

ACKNOWLEDGEMENT

I would like to thank C. Favre for his support and his direction during the writing of this article.

2. NOTATIONS AND BASICS

2.1. Basics on birational maps on surfaces. See [6, 8, 11] for details.

In this section a variety is defined over an algebraically closed field k . Recall that the resolution of singularities exists for surfaces over any algebraically closed field (see [1]).

Let X be a smooth projective surface. We denote by $N^1(X)$ the Néron-Severi group of X i.e. the group of numerical equivalence classes of divisors on X and write $N^1(X)_{\mathbb{R}} := N^1(X) \otimes \mathbb{R}$. Let $\phi : X \rightarrow Y$ be a morphism of smooth projective surfaces. It induces a natural map $\phi^* : N^1(Y)_{\mathbb{R}} \rightarrow N^1(X)_{\mathbb{R}}$. Since $\dim X = 2$, one has a perfect pairing

$$N^1(X)_{\mathbb{R}} \times N^1(X)_{\mathbb{R}} \rightarrow \mathbb{R}, \quad (\delta, \gamma) \rightarrow (\delta \cdot \gamma) \in \mathbb{R}$$

induced by the intersection form. We denote by $\phi_* : N^1(X)_{\mathbb{R}} \rightarrow N^1(Y)_{\mathbb{R}}$ the dual operator of ϕ^* .

Let X, Y be two smooth projective surfaces and $f : X \dashrightarrow Y$ be a birational map. We denote by $I(f) \subseteq X$ the indeterminacy set of f . For any curve $C \subset X$, we write $f(C) := \overline{f(C \setminus I(f))}$ the strict transformation of C .

Let $f : X \dashrightarrow X$ be a birational transformation and Γ be a desingularization of its graph. Denote by $\pi_1 : \Gamma \rightarrow X, \pi_2 : \Gamma \rightarrow X$ the natural projections. Then the diagram

$$\begin{array}{ccc} & \Gamma & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ X & \overset{f}{\dashrightarrow} & X \end{array} \quad (*)$$

is commutative and we call it a *resolution* of f .

Proposition 2.1 ([16]). *We have the following properties.*

- (i) *The morphisms π_1, π_2 are compositions of point blowups.*
- (ii) *For any point $Q \notin I(f)$, there is a Zariski open neighborhood U of p in X and an injective morphism $\sigma : U \rightarrow \Gamma$ such that $\pi_1 \circ \sigma = \text{id}$.*

Then we define the following linear maps

$$f^* = \pi_{1*} \pi_2^* : N^1(X)_{\mathbb{R}} \rightarrow N^1(X)_{\mathbb{R}},$$

and

$$f_* = \pi_{2*} \pi_1^* : N^1(X)_{\mathbb{R}} \rightarrow N^1(X)_{\mathbb{R}}.$$

Observe that $f_* = f^{-1*}$.

Note that in general we have $(f \circ g)^* \neq g^* f^*$.

For any big and nef class $\omega \in N_{\mathbb{R}}^1(X)$, we set

$$\deg_{\omega}(f) := (f^* \omega \cdot \omega),$$

the limit $\lim_{n \rightarrow \infty} \deg_{\omega}(f^n)^{1/n}$ exists and does not depend on the choice of ω (see [8, 9]). We denote this limit by $\lambda(f)$ and call it the *dynamical degree* of f .

Definition 2.2 (see [8]). Let $f : X \dashrightarrow X$ be a birational transformation on a smooth projective surface. Then f is *algebraically stable* if and only if there is no curve $V \subseteq X$ such that $f^n(V) \subseteq I(f)$ for some integer $n \geq 0$.

In the case $X = \mathbb{P}^2$, f is algebraically stable if and only if $\deg(f^n) = (\deg f)^n$ for any $n \in \mathbb{N}$.

Theorem 2.3 ([8]). *Let $f : X \dashrightarrow X$ be a birational transformation of a smooth projective surface. Then there exists a smooth projective surface $\widehat{X}$, and a proper modification $\pi : \widehat{X} \rightarrow X$ such that the lift of f to $\widehat{X}$ is an algebraically stable map.*

A *compactification* of $\mathbb{A}^2$ is a smooth projective surface X admitting a birational morphism $\pi : X \dashrightarrow \mathbb{P}^2$ that is an isomorphism above $\mathbb{A}^2 \subseteq \mathbb{P}^2$, see [11].

The following theorem which comes from [11, Proposition 2.6] and [11, Theorem 3.1] provides us a good compactification for our study.

Theorem 2.4 ([11]). *Let $f : \mathbb{A}^2 \rightarrow \mathbb{A}^2$ be a birational polynomial transformation with $\lambda(f) > 1$. Then there exists a compactification X of $\mathbb{A}^2$ satisfying the following properties.*

- (i) *The map f extends to an algebraically stable map $\widetilde{f}$ on X .*
- (ii) *There exists a $\widetilde{f}$ -fixed point $Q \in X \setminus \mathbb{A}^2$ such that $d\widetilde{f}^2|_Q = 0$.*
- (iii) *There exists an integer $n \geq 1$ such that $\widetilde{f}^n(X \setminus \mathbb{A}^2) = Q$.*

2.2. Branches of curves on surfaces. [13, 17] Let X be a smooth projective surface over an algebraically closed field k , C be an irreducible curve in X and p be a point in C .

Let $I_{C,p}$ be the prime ideal associate to C in the local function ring $O_{X,p}$ and $\widehat{I_{C,p}}$ be the completion of $I_{C,p}$ in the completions of local function ring $\widehat{O_{X,p}}$.

Definition 2.5. A *branch* of C at p is a prime ideal $\mathfrak{p}_{C,p}$ of $\widehat{O_{X,p}}$ with height 1 which contains $\widehat{I_{C,p}}$.

Let $i : \tilde{C} \rightarrow C$ is a normalization of C and $\tilde{p}$ a point in $i^{-1}(p)$, the morphism $i : \tilde{C} \rightarrow C$ induces a morphism $i^* : \widehat{\mathcal{O}_{X,p}} \rightarrow \widehat{\mathcal{O}_{\tilde{C},\tilde{p}}}$ between the completions of local function rings. The map $\tilde{p} \mapsto \mathfrak{p}_{\tilde{p}} := \ker i^*$ give us an one to one correspondents between the set of branches of C at p and the set $i^{-1}(p)$. So we may view a branch of C as a triple $(i, \tilde{C}, \tilde{p})$ up to the following equivalent relations: two branches $(i, \tilde{C}, \tilde{p})$ and $(i', \tilde{C}', \tilde{p}')$ are equivalent if there exists an isomorphism $\theta : \tilde{C} \rightarrow \tilde{C}'$ over C sending $\tilde{p}$ to $\tilde{p}'$.

Given any two different branches $C_p = (i, \tilde{C}, \tilde{p})$ and $D_q = (j, \tilde{D}, \tilde{q})$ at p and q respectively, we define the intersection number by

$$(C_p \cdot D_q)_r = \begin{cases} \dim_k \widehat{\mathcal{O}_{X,p}} / (\mathfrak{p}_{C,\tilde{p}} + \mathfrak{p}_{D,\tilde{p}}), & \text{when } p = q = r; \\ 0, & \text{ohterwise.} \end{cases}$$

If $D := j(\tilde{D})$ is different from $C := i(\tilde{C})$, we may also define the intersection numbers by

$$(C_p \cdot D)_r = (D \cdot C_p)_r = \sum_{\tilde{q} \in j^{-1}(r)} (C_r \cdot (j, \tilde{D}, \tilde{q}))$$

and

$$(C \cdot D)_r = \sum_{\tilde{p} \in i^{-1}(r), \tilde{q} \in j^{-1}(r)} ((i, \tilde{C}, \tilde{p}) \cdot (j, \tilde{D}, \tilde{q})).$$

The following proposition shows that our definition is agree with the usual definition of intersection number of curves.

Proposition 2.6. *We have*

$$(C \cdot D) = \sum_{\tilde{p} \in \tilde{C}, \tilde{q} \in \tilde{D}, r \in X} ((i, \tilde{C}, \tilde{p}) \cdot (j, \tilde{D}, \tilde{q}))_r = \sum_{r \in C \cap D} \sum_{\tilde{p} \in i^{-1}(r), \tilde{q} \in j^{-1}(r)} ((i, \tilde{C}, \tilde{p}) \cdot (j, \tilde{D}, \tilde{q}))_r.$$

Let Z be a smooth projective surface and $f : X \dashrightarrow Z$ be a birational map. If f does not contract C then we denote by $f(C_p) := (f \circ i, \tilde{C}, \tilde{p})$ and call it the strict transformation of C_p . Observe that $f(C_p)$ is a branch of $f(C)$ and when $p \notin I(f)$, we have that $f(C_p)$ is a branch of curve at $f(p)$.

If f is regular at p , we write

$$f_* C_p = \begin{cases} f(C_p), & \text{when } f \text{ does not contracts } C_p; \\ 0, & \text{ohterwise.} \end{cases}$$

Let Y be another smooth projective surface and $\pi : Y \rightarrow X$ be a birational morphism. Denote by $\pi^\# C_p := \pi^{-1}(C)$ the strict transformation of C_p . Let E_i , $i = 1, \dots, m$ be the exceptional curves of π . There is a unique sequence of non negative integers $(a_i)_{0 \leq i \leq m}$ such that for any irreducible curve D in Y different from $\pi^\# C$, we have $(C_p \cdot \pi_* D) = (\pi^\# C_p + \sum_{i=1}^m a_i E_i \cdot D)$. Denote by $\pi^* C_p := \pi^\# C_p + \sum_{i=1}^m a_i E_i$ and call it the pull back of C_p .

Proposition 2.7. *We have the following properties.*

- (i) *We have $\pi_* \pi^* C_p = C_p$.*

- (ii) For any irreducible curve (resp. branch of curve) D in Y different from $\pi^\#C$ (resp. $\pi^\#C_p$), we have

$$(\pi^*C_p \cdot D) = (C_p \cdot \pi_*D).$$

- (iii) For any curve (resp. branch of curve) D in X different from C (resp. C_p) then we have

$$(C_p \cdot D) = \sum_{x \in \pi^{-1}(p)} (\pi^\#C_p \cdot \pi^*D)_x.$$

2.3. Absolute values on fields. (See [20]) At first, we recall the definition of absolute values on fields.

Definition 2.8. An absolute value on a field K is a function

$$|\cdot| : K \rightarrow [0, +\infty)$$

satisfies the following properties:

- (i) $|x| = 0$ if and only if $x = 0$;
- (ii) $|xy| = |x||y|$;
- (iii) $|x + y| \leq |x| + |y|$.

The absolute value is said to be nonarchimedean if it satisfies (iii') $|x + y| \leq \max\{|x|, |y|\}$.

In our paper, a key idea is to use all nonequivalent absolute values on a field at the same time to get more information. For completeness, we give a brief introduction of *place* on number fields and function fields of curves, see [20] for more discussions.

Definition 2.9. The set of places on $\mathbb{Q}$, which is denoted by $\mathcal{M}_{\mathbb{Q}}$ consists of the following absolute values:

- (i) the usual absolute value $|\cdot|_{\infty} : x \mapsto \max\{x, -x\}$ for any $x \in \mathbb{Q}$;
- (ii) for any prime p , the p -adic absolute value $|\cdot|_p : |p^n a/b|_p \mapsto n$ where $a, b \in \mathbb{Z}$ and $p \nmid ab$.

The usual absolute value $|\cdot|_{\infty}$ is the only one archimedean place of $\mathbb{Q}$.

Let $K/\mathbb{Q}$ be a number field. The set of places on K is denoted by $\mathcal{M}_K$ and consists of all absolute values on K whose restriction to $\mathbb{Q}$ is one of the places in $\mathcal{M}_{\mathbb{Q}}$. Further we denote by $\mathcal{M}_K^{\infty} := \{v \in \mathcal{M}_K \mid |\cdot|_v|_{\mathbb{Q}} = |\cdot|_{\infty}\}$ the set of archimedean places; and by $\mathcal{M}_K^0 := \{v \in \mathcal{M}_K \mid |\cdot|_v|_{\mathbb{Q}} = |\cdot|_p \text{ for some prime } p \in \mathbb{N}\}$ the set of nonarchimedean places.

When v is non-archimedean, there exists an embedding $\sigma_v : K \hookrightarrow \mathbb{C}$ or $\mathbb{R}$ such that $|\cdot|_v$ is the restriction of the usual absolute value.

Definition 2.10. For any place $v \in \mathcal{M}_K$, we denote by K_v the completion of K at v and by $n_v := [K_v : \mathbb{Q}_v]$ the *local degree* of v .

In parallel, we define the set of places on function fields.

Definition 2.11. Let C be a smooth projective curve defined over an algebraically closed field k and $L := K(C)$ be the function field of C . The set of places on L , denoted by $\mathcal{M}_L$ consists of the absolute values of forms: for any closed point $p \in C$,

$$|\cdot|_p : x \mapsto e^{\text{ord}_p(x)}$$

for any $x \in L$.

Let K/L be a finite field extension. The set of places on K is denoted by $\mathcal{M}_K$ and consists of all absolute values on K whose restriction to L is one of the places in $\mathcal{M}_L$. In this case, all the places in $\mathcal{M}_K$ are nonarchimedean. We define $\mathcal{M}_K^0 = \mathcal{M}_K$ and $\mathcal{M}_K^\infty = \emptyset$ for convenience. For any place $v \in \mathcal{M}_K$, we denote by $n_v := [K_v : L_v]$ the local degree of v .

Let K/L be a finite field extension where $L = \mathbb{Q}$ or a function field $K(C)$ of a curve C . We denote by $O_v := \{x \in K \mid |x|_v \leq 1\}$ for any $v \in \mathcal{M}_K$. In the number field case, we denote by O_K the ring of integers of K i.e. $O_K = \{x \in K \mid |x|_v \leq 1 \text{ for all } v \in \mathcal{M}_K^0\}$.

Proposition 2.12 (Product Formula). *Let K/L be a finite field extension where $L = \mathbb{Q}$ or a function field $K(C)$ of a curve C . Then we have*

$$\prod_{v \in \mathcal{M}_K} |x|_v^{n_v} = 1$$

for all $x \in K^*$.

2.4. Metrics on projective varieties defined over a valued field. A field with an absolute value is called a valued field.

At first, we define a metric on the projective space over any valued field as follows.

Definition 2.13. Let $(K, |\cdot|_v)$ be a valued field. For any integer $n \geq 1$, we define a metric d_v on the projective space $\mathbb{P}^n(K)$ by

$$d_v([x_0 : \cdots : x_n], [y_0 : \cdots : y_n]) = \frac{\max_{0 \leq i, j \leq n} |x_i y_j - x_j y_i|_v}{\max_{0 \leq i \leq n} |x_i|_v \max_{0 \leq j \leq n} |y_j|_v}$$

for any two points $[x_0 : \cdots : x_n], [y_0 : \cdots : y_n] \in \mathbb{P}^n(K)$.

Observe that when $|\cdot|_v$ is archimedean, the metric d_v is not induced by a smooth riemannian metric. However it is equivalent to the restriction of Fubini-Study metric on $\mathbb{P}^n(\mathbb{C})$ or $\mathbb{P}^n(\mathbb{R})$ to $\mathbb{P}^n(K)$ induced by σ_v .

More generally, for a projective variety X defined over $(K, |\cdot|_v)$, if we fix an embedding $\iota : X \hookrightarrow \mathbb{P}^n(K)$, we may restrict the metric d_v on $\mathbb{P}^n(K)$ to a metric $d_{v, \iota}$ on X . This metric depends on the choice of embedding ι in general, but for different embeddings ι_1 and ι_2 , the metrics d_{v, ι_1} and d_{v, ι_2} are equivalent. Since we are mostly intersecting in the topology induced by the metric we shall usually write d_v for $d_{v, \iota}$ for simplicity.

3. A CRITERION FOR A CURVE TO BE PERIODIC

Our aim in this section is to prove Theorem 1.2 from the introduction. Let us recall the setting:

- (i) Let X be a smooth projective surface over an algebraically closed field.
- (ii) Let $f : X \dashrightarrow X$ be an algebraically stable birational transformation.
- (iii) Let C be an irreducible curve in X such that f^n does not contract C and $f^n(C) \cap I(f) \neq \emptyset$ for all n .

Let us begin with the following special case.

Lemma 3.1. *Let x be a point in $I(f) \cap C$. If there is a branch s of C at x such that $f^n(s)$ is again a branch at x for all $n \geq 0$, then C is fixed.*

Proof of Lemma 3.1. Since f is birational, we chose a resolution of f as in the diagram (*) in Sectin 2.1.

If C is not fixed, we have $f(s) \neq s$ and set $A = (s \cdot f(s))_x < \infty$. By Proposition 2.1, π_2 is invertible on a Zariski neighbourhood of x . Set F_x the fiber of π_1 at x .

For any $m \geq 0$, we have,

$$\begin{aligned}
 ((f^m(s) \cdot f^{m+1}(s))_x &= \sum_{y \in F_x} (\pi_1^\# f^m(s) \cdot \pi_1^* f^{m+1}(s))_y \\
 &\geq (\pi_1^\# f^m(s) \cdot \pi_1^* f^{m+1}(s))_{\pi_2^{-1}(x)} \\
 &= (\pi_1^\# f^m(s) \cdot \pi_1^\# f^{m+1}(s))_{\pi_2^{-1}(x)} + (\pi_1^\# f^m(s) \cdot F_x)_{\pi_2^{-1}(x)} \\
 &= (f^{m+1}(s) \cdot f^{m+2}(s))_x + (\pi_1^\# f^m(s) \cdot F_x)_{\pi_2^{-1}(x)} \\
 &\geq (f^{m+1}(s) \cdot f^{m+2}(s))_x + 1.
 \end{aligned}$$

It follows that $A = (s \cdot f(s))_x \geq (f^m(s) \cdot f^{m+1}(s))_x + m \geq m$ for all $m \geq 0$ which yield a contradiction. $\square$

We now treat the general case.

Proof of Theorem 1.2. Recall that f^n does not contract C and $f^n(C) \cap I(f) \neq \emptyset$ for all n . By Lemma 3.1, it is sufficient to find a point $x \in I(f) \cap C$ such that the image by f^n of the branch of C at x is again a branch of curve at x for all $n \geq 0$. By contradiction we suppose that C is not periodic.

To do so, we introduce

$$P(f) = \{x \in I(f) \mid \text{there is } n_1 > n_2 \geq 0 \text{ such that } f^{-n_1}(x) = f^{-n_2}(x)\}$$

and

$$O(f) = \{f^{-n}(x) \mid x \in P(f) \text{ and } n \geq 0\}.$$

By definition, $O(f)$ is finite. Since f is algebraically stable, $O(f) = O(f^n)$ for all $n \geq 1$. Replacing f by f^l for a suitable $l \geq 1$, we may assume that $O(f) = P(f)$. Set $N(f) = I(f) - P(f)$.

At first, we prove

Lemma 3.2. *For all $n \geq 0$, $f^n(C) \cap O(f) \neq \emptyset$.*

Proof of Lemma 3.2. We assume that $I(f) = \{p_1, \dots, p_m\}$ and define the map

$$F = (f^{-1}, \dots, f^{-1}) : X^m \dashrightarrow X^m.$$

Denote by π_i the projection to the i -th factor and set

$$D = \bigcup_{i=1}^m \pi_i^{-1}(C).$$

Set $q = (p_1, \dots, p_m)$ a point in X^m . Our hypothesis implies that $F^n(q) \in D$ for all $n \geq 0$. Let Z' be the Zariski closure of $\{F^n(q) | n \geq 0\}$. Then $Z' \subseteq D$. Let Z be the union of all positively dimensional irreducible components of Z' . If $Z = \emptyset$, then p_i is f^{-1} -preperiodic for all i . Otherwise since $\{F^n(q) | n \geq 0\} \cap I(F) = \emptyset$, the proper transformation of Z by F is well defined and satisfies $F(Z) = Z$, hence each irreducible component of Z is periodic. Let l be a common period for all components of Z . For any component W , there exists one $i \in \{1, \dots, m\}$ such that $W \subseteq \pi_i^{-1}(C)$. In other words, there exists $k \geq 0$ such that $f^{-ln-k}(p_i) \in C$ for all $n \geq 0$. If p_i is not f^{-1} -preperiodic, then C is the Zariski closure of $\{f^{-ln}(p_i) | n \geq 0\}$ which is f^{-l} -invariant. This implies C to be periodic and contradicts to our hypothesis. It follows that p_i is f^{-1} -preperiodic.

Repeating the same argument for $f^n(C)$, we have $f^n(C) \cap O(f) \neq \emptyset$ for all $n \geq 0$. $\square$

Denote by $D(n)$ the number of branches of $f^n(C)$ at points of $O(f)$ and set

$$S = \{x \in O(f) | \text{there are infinitely many } n \geq 0 \text{ such that } x \in f^n(C)\}.$$

Since $f^{-1}(O(f)) \subseteq O(f)$, we have $f^n(x) \in O(f)$ implies $x \in O(f)$. It follows that $D(n)$ is decrease and nonzero. There is an integer M , such that for $n \geq M$, $D(n)$ stabiles. We may replace C by $f^M(C)$ and assume that $D(n)$ is constant for $n \geq 0$. Then for each branch of curve of $f^n(C)$ at a point in $O(f)$, its image by f is a branch of $f^{n+1}(C)$ at a point of $O(f)$. By the definition of S and the finiteness of $O(f)$, we have

$$f^n(C) \cap O(f) = f^n(C) \cap S$$

for all integer $n \geq 0$.

According to Lemma 3.1, the following lemma concludes our proof.

Lemma 3.3. *Replacing f by a positive iterate, there exists a point $x \in C \cap S$ for which there is a branch s of C such that $f^n(s)$ is again a branch of curve at x for all $n \geq 0$.*

$\square$

Proof of Lemma 3.3. Chose a resolution of f as in the diagram (*) in Section 2.1. For any point $x \in S$, denote by F_x the fibre of π_1 at x and $E_x = \pi_2(F_x) \cap S$.

We have $E_x \neq \emptyset$. Otherwise, there exists $n \geq 0$ for which $x \in f^n(C)$ and a branch s of $f^n(C)$ at x . The assumption $E_x = \emptyset$ implies that $f(s)$ is not a branch at any point in S . This shows that $D(n+1) < D(n)$ and we get a contradiction.

On the other hand, let x_1, x_2 be two different points in S . If $E_{x_1} \cap E_{x_2} \neq \emptyset$, there exists $y \in S$ such that $y \in \pi_2(F_{x_1}) \cap \pi_2(F_{x_2})$. By Zariski's main theorem, $\pi_2^{-1}(y)$ is a connected curve meeting F_{x_1} and F_{x_2} . So $\pi_1(\pi_2^{-1}(y))$ is a curve and it is contracted by f to $y \in S \subseteq I(f)$. This contradicts the fact that f is algebraically stable. So we have

$$E_{x_1} \cap E_{x_2} = \pi_2(F_{x_1}) \cap \pi_2(F_{x_2}) \cap S = \emptyset.$$

Set $T = \coprod_{x \in S} E_x$ a subset of S . Since $\#E_x \geq 1$ for all x , we have $\#T \geq \#S$. It follows that $T = S$ and $\#E_x = 1$ for all $x \in S$. This allows us to define a map $G : S \rightarrow S$ sending $x \in S$ to the unique point in E_x . Then G is an one to one map. For all $n \geq M$, f sends a branch of $f^n(C)$ at a point $x \in S$ to a branch of $f^{n+1}(C)$ at the point $G(x)$. By replacing f by $f^{(\#S)!}$, we may assume that $G = \text{id}$. Then for any $x \in S \cap C$ and s a branch of C at x , we have $f^n(s)$ is again a branch at x for all $n \geq 0$. $\square$

4. THE DML PROPERTY

Definition 4.1. Let X be a smooth surface defined over an algebraically closed field, and $f : X \dashrightarrow X$ be a birational transformation. We say that the pair (X, f) satisfies the property DML if for any curve C on X and for any closed point $p \in X$ such that $f^n(p) \notin I(f)$ for all $n \geq 0$, the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is a union of at most finitely many arithmetic progressions.

In our setting the property DML is infact equivalent to the following seemingly stronger property.

Proposition 4.2. *Let X be a smooth surface defined over an algebraically closed field, and $f : X \dashrightarrow X$ be a birational transformation. The following statements are equivalent.*

- (1) *The pair (X, f) satisfies the property DML.*
- (2) *For any curve C on X and any closed point $p \in X$ such that $f^n(p) \notin I(f)$ for all $n \geq 0$ and the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite, then p is preperiodic or C is periodic.*

Proof. Suppose (1) holds. Let C be any curve in X and p be a closed point in X such that $f^n(p) \notin I(f)$ for all $n \geq 0$. Assume that the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite. The property DML of (X, f) implies that there are integers $a > 0$ and $b \geq 0$ such that $f^{an+b}(p) \in C$ for all $n \geq 0$. If p is not preperiodic, the set $O_{a,b} := \{f^{an+b}(p) \mid n \geq 0\}$ is Zariski dense in C and $f^a(O_{a,b}) \subseteq O_{a,b}$. It follows that $f^a(C) \subseteq C$, hence C is periodic.

Suppose (2) holds. If the set $S := \{n \in \mathbb{N} \mid f^n(p) \in C\}$ is finite or p is preperiodic, there is nothing to prove. So we assume that S is infinite and p is not preperiodic. The property (2) implies that C is periodic. There exists an integer $a > 0$ such that $f^a(C) \subseteq C$. So S is a finite union of sets of form $\{an + b \mid n \geq 0\}$ where $b \geq 0$ is an integer, and it follows that (X, f) satisfies the property DML. $\square$

Theorem 4.3. *Let X be a smooth surface defined over an algebraically closed field, and $f : X \dashrightarrow X$ be a birational transformation, then the following properties hold.*

- (i) *For any $m \geq 1$, (X, f) satisfies the property DML if and only if (X, f^m) satisfies the property DML.*
- (ii) *Suppose U is an open subset of X such that the restriction $f|_U : U \rightarrow U$ is a morphism. Then (X, f) satisfies the property DML, if and only if $(U, f|_U)$ satisfies the property DML.*
- (iii) *Suppose $\pi : X \rightarrow X'$ is a birational morphism between smooth projective surfaces, and $f : X \dashrightarrow X$, $f' : X' \dashrightarrow X'$ are birational maps such that $\pi \circ f = f' \circ \pi$. If the pair (X, f) satisfies the property DML, then (X', f') satisfies the property DML.*
- (iv) *Suppose $\pi : X \rightarrow X'$ is a birational morphism between smooth projective surfaces, and $f : X \dashrightarrow X$, $f' : X' \dashrightarrow X'$ are birational transformations such that $\pi \circ f = f' \circ \pi$. If f' is algebraically stable and the pair (X', f') satisfies the property DML, then (X, f) satisfies the property DML.*

Definition 4.4. Let X be a smooth projective surface defined over an algebraically closed field and $f : X \dashrightarrow X$ be a birational transformation. We say that (X', f') is a birational model of (X, f) if there is a birational map $\pi : X' \dashrightarrow X$ such that

$$f' = \pi^{-1} \circ f \circ \pi.$$

Corollary 4.5. *Let X be a smooth projective surface defined over an algebraically closed field and $f : X \dashrightarrow X$ be an algebraically stable birational transformation such that (X, f) satisfies the property DML. Then all birational models (X', f') of (X, f) satisfy the property DML.*

Proof of Theorem 4.3. (i). The "only if" part is trivial, so we only have to due with the "if" part. We assume that (X, f^m) satisfies the property DML. Let C be a curve in X and p be a point in X such that $f^n(p) \notin I(f)$ for all $n \geq 0$ and the set

$$\{n \in \mathbb{N} \mid f^n(p) \in C\}$$

is infinite. Since

$$\{n \in \mathbb{N} \mid f^n(p) \in C\} = \bigcup_{i=0}^{m-1} \{n \in \mathbb{N} \mid f^{nm}(f^i(p)) \in C\},$$

then for some i , the set $\{n \in \mathbb{N} \mid f^{nm}(f^i(p)) \in C\}$ is infinite. Since (X, f^m) satisfies the property DML, C is periodic or $f^i(p)$ is preperiodic. It follows that C is periodic or $f(p)$ is preperiodic.

(ii). If (X, f) satisfies the property DML, since $f|_U : U \rightarrow U$ is a morphism, $(U, f|_U)$ satisfies the property DML.

Conversely suppose that $(U, f|_U)$ satisfies the property DML. Let C be an irreducible curve in X and p be a closed point in X such that $f^n(p) \notin I(f)$ for all $n \geq 0$ and the set

$$\{n \in \mathbb{N} \mid f^n(p) \in C\}$$

is infinite. The set $E = X - U$ is a proper closed subvariety of X . If $p \in U$, then we have that $C \not\subseteq E$. Since $(U, f|_U)$ satisfies the property DML, we have either p is preperiodic or C is periodic. Otherwise, we may assume that for all $n \geq 0$, $f^n(p) \in E$, then the Zariski closure D of $\{f^n(p) \mid n \geq 0\}$, contains in E . We assume that p is not preperiodic, then $C \subseteq D$. Since D is fixed, we have that C is periodic.

(iii). It is sufficient to treat the case when π is the blowup at a point $q \in X'$. Let C' be a curve in X' and p' be a point in X' such that $f'^n(p') \notin I(f')$ for all $n \geq 0$ and the set $\{n \in \mathbb{N} \mid f'^n(p') \in C'\}$ is infinite. We assume that p' is not a periodic point, and then we have that for n large enough, $f'^n(p') \neq q$. Replacing p by $f'^m(p')$ for some m large enough, we assume that $f'^n(p') \neq q$ for all $n \geq 0$. Set $p = \pi^{-1}(p')$ and $C = \pi^{-1}(C')$, then we have $f^n(p) \notin I(f)$ for all $n \geq 0$ and the set

$$\{n \in \mathbb{N} \mid f^n(p) \in C\}$$

is infinite. This implies C and then C' to be periodic.

(iv). Let $C \subseteq X$ be a curve and p be a point in X such that $f^n(p) \notin I(f)$ for all $n \geq 0$ and the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite. We may assume that C is irreducible. Let E be the exceptional locus of π .

Lemma 4.6. *If $C \subseteq E$ and $\pi(C)$ is a point in $I(f')$, then (iv) holds.*

Proof of Lemma 4.6. Set $q := \pi(C) \in I(f')$. Since f' is algebraically stable, we have $q \notin I((f')^{-n})$ and

$$\pi(f^{-n}(C)) = (f')^{-n}(q)$$

for all $n \geq 1$. It follows that $f^{-n}(C)$ is a point or an exceptional curve of π for $n \geq 1$.

If there exists $l \geq 1$ such that $f^{-l}(C)$ is a point, we pick two integers $n_1 > n_2 \geq l$ such that $f^{n_1}(p), f^{n_2}(p) \in C$. Then $f^{n_1-l}(p) = f^{n_2-l}(p)$, which implies p to be preperiodic.

Otherwise $f^{-n}(C)$ is an exceptional curve of π , for all $n \geq 0$. Since there are only finitely many irreducible components of E , we have that C is periodic. $\square$

Denote by $K = \pi^{-1}(I(f'))$.

Lemma 4.7. *If there are infinitely many $n \geq 0$ such that $f^n(p) \in K$, then (iv) holds.*

Proof of 4.7. There is an irreducible component F of K such that the set $\{n \geq 0 \mid f^n(p) \in F\}$ is infinite.

If F is a point, then p is preperiodic.

Otherwise F is a curve, then $F \subseteq E$ and $\pi(F) \subseteq I(f')$. Suppose that p is not preperiodic, Lemma 4.6 shows that F is periodic. Then $F' = \bigcup_{k \geq 0} f^k(F)$ is a curve and $f^n(p) \subseteq F'$ for all $n \geq 0$. If $C \subseteq F'$, then C is periodic. If $C \not\subseteq F'$, then $C \cap F'$ is finite, and this shows that p is preperiodic. $\square$

Lemma 4.8. *If $C \subseteq E$, then (iv) holds.*

Proof. By Lemma 4.7, we may assume that there exists an integer $N \geq 0$, such that $f^n(p) \notin K$ for all $n \geq N$.

Set $q := \pi(C)$. By Lemma 4.6, we assume that $q \notin I(f')$. Then we have

$$\pi(f^{N+l}(p)) = f'^l(\pi(f^N(p)))$$

for $l \geq 0$. It follows that there are infinitely many $l \geq 0$, such that $f'^l(\pi(f^N(p))) = q$. Then q is preperiodic and the orbit of $f'^N(q)$ does not meet $I(f')$. Since $\pi(f^n(C)) = f'^n(q)$ for all $n \geq 0$, we have $f^n(C) \subseteq \bigcup_{k \geq N} \pi^{-1}(f^k(q))$ for all $n \geq N$. Hence either C is periodic or for some $n \geq 1$, $f^n(C)$ is a point. In the second case, we conclude that p is preperiodic. $\square$

Let $L = K \cup E$.

Lemma 4.9. *If there are infinitely many $n \geq 0$ such that $f^n(p) \in L$, then (iv) holds.*

Proof of Lemma 4.9. There is an irreducible component F such that $\{n \geq 0 \mid f^n(p) \in F\}$ is infinite.

If F is a point, then p is preperiodic.

Otherwise F is a curve, then $F \subseteq E$. Suppose that p is not preperiodic, Lemma 4.8 shows that F is periodic. Then $F' = \bigcup_{k \geq 0} f^k(F)$ is a curve and $f^n(p) \in F'$ for all $n \geq 0$. If $C \subseteq F'$, then C is periodic. Otherwise $C \not\subseteq F'$, we have that $C \cap F'$ is finite and then p is preperiodic. $\square$

We may assume that there is an integer $M \geq 0$, such that $f^n(p) \notin L$ for all $n \geq M$.

If $C \not\subseteq E$, $\pi(C)$ is a curve. For all $l \geq 0$ we have

$$\pi(f^{M+l}(p)) = f'^l(\pi(f^M(p))) \notin I(f').$$

Since (X', f') satisfies the property DML, either $\pi(C)$ is periodic or $\pi(p)$ is preperiodic. When $\pi(C)$ is periodic, we have C is periodic. Otherwise $\pi(p)$ is preperiodic. For any $l \geq 0$, π is invertible on some Zariski neighborhood of the point $f'^l(\pi(f^M(p)))$ and then we conclude that p is preperiodic. $\square$

Proof of Corollary 4.5. Pick Y a desingularization of the graph of f and set π_1, π_2 the projections which make the diagram

$$\begin{array}{ccc} & Y & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ X & \xrightarrow{\phi} & X' \end{array}$$

to be commutative. Since f is algebraically stable, its lift to Y satisfies the property DML by Theorem 4.3 (iv). We conclude that (X', f') satisfies the property DML by Theorem 4.3 (iii). $\square$

5. LOCAL DYNAMICAL MORDELL LANG THEOREM

The aim of this section is to prove Theorem 1.1.

Proof of Theorem 1.1. Pick a resolution of f as the diagram (*) in Sectin 2.1 and recall Proposition 2.1. Assume that for all $n \geq 0$, $f^n(p) \neq Q$. There is an infinite sequence $\{n_k\}_{k \geq 0}$ such that $f^{n_k}(p) \in C \setminus \{Q\}$. It follows that $f^{n_k-m}(p) \in f^{-m}(C)$ for k large enough. With $k \rightarrow \infty$, we get $Q \in f^{-m}(C)$ for all $m \geq 0$.

If $C \neq f^{-1}(C)$, then we have $f^{-m}(C) \neq f^{-m-1}(C)$ for all $m \geq 0$. We may calculate the local intersection of them at Q . We have

$$\begin{aligned} (f^{-m}(C) \cdot f^{-m-1}(C))_Q &= \sum_{x \in \pi_2^{-1}(Q)} (\pi_2^* f^{-m}(C) \cdot \pi_2^\# f^{-m-1}(C))_x \\ &= \sum_{x \in \pi_2^{-1}(Q)} ((\pi_2^\# f^{-m}(C) + \sum_{i=1}^s v_{E_i}(f^{-m}(C))E_i) \cdot \pi_2^\# f^{-m-1}(C))_x \end{aligned} \quad (5.1)$$

where $E_i, 1 \leq i \leq s$ are irreducible exceptional curves of π_2 . Since

$$\text{Supp}(\sum_{i=1}^s v_{E_i}(f^{-m}(C))E_i) = \bigcup_{1 \leq i \leq s} E_i = \pi_2^{-1}(Q),$$

we have

$$\begin{aligned} (5.1) &= \sum_{x \in \pi_2^{-1}(Q)} (\pi_2^\# f^{-m}(C) \cdot \pi_2^\# f^{-m-1}(C))_x + ((\sum_{i=1}^s v_{E_i}(f^{-m}(C))E_i) \cdot \pi_2^\# f^{-m-1}(C)) \\ &\geq \sum_{x \in \pi_2^{-1}(Q)} (\pi_2^\# f^{-m}(C) \cdot \pi_2^\# f^{-m-1}(C))_x + 1 \\ &= (\sigma(f^{-m-1}(C)) \cdot \sigma(f^{-m-2}(C)))_{\sigma(Q)} + 1 \\ &= (f^{-m-1}(C) \cdot f^{-m-2}(C))_Q + 1. \end{aligned}$$

It follows that

$$0 < (f^{-m}(C) \cdot f^{-m-1}(C))_Q \leq (f^{-m+1}(C) \cdot f^{-m}(C))_Q - 1 \leq \dots \leq (C \cdot f^{-1}(C))_Q - m$$

for all $m \geq 0$, which yields a contradiction. So we have $C = f^{-1}(C)$ and then $f(C) = C$. $\square$

Observe that our proof of Theorem 1.1 actually gives

Proposition 5.1. *Let X be a projective surface over an algebraically closed field and $f : X \dashrightarrow X$ be a birational map with a fixed point $Q \in I(f^{-1}) \setminus I(f)$. Then all periodic curves passing through Q are fixed.*

6. THE CASE $\lambda(f) = 1$

In this section, we prove Theorem A in the case $\lambda(f) = 1$. Denote by K an algebraically closed field of characteristic 0.

Recall from [8] and [11], that if $\lambda(f) = 1$, then we are in one of the following two cases:

- (1) there exists a smooth projective surface X and an automorphism f' on X such that the pair (X, f') is birationally conjugates to $(\mathbb{A}^2, f)$;
- (2) in suitable affine coordinate, $f(x, y) = (ax + b, A(x)y + B(x))$ where A and B are polynomials with $A \neq 0$ and $a \in K^*, b \in K$.

In case (1), [3, Theorem 1.3] of Bell, Ghioca and Tucker implies our Theorem A immediately, so we only have to consider case (2). Though out this section we shall always assume that

$$f(x, y) = (ax + b, A(x)y + B(x)) \quad (**)$$

with $A, B \in K[x]$, $A \neq 0$, $a \in K^*$ and $b \in K$.

6.1. Algebraically stable models. Any map of the form $(**)$ can be made to be algebraically stable in a suitable Hirzebruch surface $\mathbb{F}_n$ for some $n \geq 0$. It is convenient to work with the presentation of these surfaces as a quotient by $(\mathbb{G}_m)^2$, as in [2]. By definition, the set of closed point $\mathbb{F}_n(K)$ is the quotient of $\mathbb{A}^4(K) \setminus (\{x_1 = 0 \text{ and } x_2 = 0\} \cup \{x_3 = 0 \text{ and } x_4 = 0\})$ by the equivalence relation generated by

$$(x_1, x_2, x_3, x_4) \sim (\lambda x_1, \lambda x_2, \mu x_3, \mu/\lambda^n x_4)$$

for $\lambda, \mu \in K^*$. We denote by $[x_1, x_2, x_3, x_4]$ the equivalence class of (x_1, x_2, x_3, x_4) . We have a natural morphism $\pi_n : \mathbb{F}_n \rightarrow \mathbb{P}^1$ given by $\pi_n([x_1, x_2, x_3, x_4]) = [x_1 : x_2]$ which makes $\mathbb{F}_n$ into be a locally trivial $\mathbb{P}^1$ fibration.

We shall look at the embedding

$$i_n : \mathbb{A}^2 \hookrightarrow \mathbb{F}_n : (x, y) \mapsto [x, 1, y, 1].$$

Then $\mathbb{F}_n \setminus \mathbb{A}^2$ is union of two lines: one is the fiber at infinity F_∞ of π_n , and the other one is a section which we denote by L_∞ .

Recall that f has form $(**)$. For each $n \geq 0$, set $d = \max\{\deg A, \deg B - n\}$. By the embedding i_n , the map f extends to a birational transformation

$$f_n : [x_1, x_2, x_3, x_4] \mapsto [ax_1 + bx_2, x_2, A(x_1/x_2)x_2^d x_3 + B(x_1/x_2)x_2^{d+n} x_4, x_2^d x_4]$$

on $\mathbb{F}_n$. For any $n \geq \deg B - \deg A + 1$, we have $d = \deg A$ and

$$I(f_n) = \{[x_1, x_2, x_3, x_4] \in \mathbb{F}_n \mid x_2 = x_3 = 0\}.$$

The unique curve which is contracted by f_n is $F_\infty = \{x_2 = 0\}$ and its image is $f_n(F_\infty) = [1, 0, 1, 0]$. It implies the following:

Proposition 6.1. *For $n \geq \deg B - \deg A + 1$, f_n is algebraically stable on $\mathbb{F}_n$ and contracts the curve F_∞ to the point $[1, 0, 1, 0]$.*

6.2. The attracting case. In the remaining of this section, we shall fix an integer m such that the extension of f to $\mathbb{F}_m$ is algebraically stable. For simplicity, we write f for the map f_m induced by f on $\mathbb{F}_m$.

Proposition 6.2. *Let $|\cdot|$ be an absolute value on K such that $|a| > 1$. Then $(\mathbb{F}_m, f)$ satisfies the property DML.*

Proof. Since $a \neq 1$, by changing coordinate, we may assume that $f = (ax, A(x)y + B(x))$. Since f contracts the fiber at infinity F_∞ to $O := L_\infty \cap F_\infty$, the point O is fixed and the two eigenvalues of df at O are $1/a$ and 0 . Since $|a| > 1$, there is a neighbourhood U of O , such that $U \cap I(f) = \emptyset$, $f(U) \subseteq U$ and $f^n \rightarrow O$ uniformly on U .

Let C be an irreducible curve in $\mathbb{P}_K^2$ and p be a point in $\mathbb{A}_K^2$ such that the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite. By Lemma 4.3 we may assume that $p \in \mathbb{A}_K^2$ and $C \not\subseteq L_\infty \cup F_\infty$.

If $C \cap F_\infty = \{O\}$, there is an open set V of $\mathbb{P}_K^1$, such that $[1 : 0] \in V$ and $\pi_m^{-1}(V) \cap C \subseteq U$. Since $|a| > 1$, for n large enough, $f^n(p) \in \pi_m^{-1}(V)$. So there is an integer $n_1 > 0$ such that $f^{n_1}(p) \in U$. Theorem 1.1 implies that the curve C is fixed.

We may assume now that $f^n(C) \cap F_\infty \neq \{O\}$ for all $n \geq 0$.

If $C \cap F_\infty = \emptyset$, then C is a fiber of the rational fibration $\pi_m : \mathbb{F}_m \rightarrow \mathbb{P}^1$. Since $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite, the curve C is fixed.

Finally assume that $f^n(C) \cap F_\infty \neq \emptyset$ for all $n \geq 0$. Since f contracts F_∞ to O , we have

$$f^n(C) \cap I(f) \neq \emptyset,$$

and we conclude by Theorem 1.2 that C is periodic in this case. $\square$

6.3. The general case.

Proposition 6.3. *The pair $(\mathbb{F}_m, f)$ satisfies the property DML.*

Proof. Let C be a curve in $\mathbb{F}_m$, and p be a point in $\mathbb{A}_K^2$ such that the set $\{n \geq 0 \mid f^n(p) \in C\}$ is infinite. We may assume that the transcendence degree of K is finite, since we can find a subfield of K such that it has finite transcendence degree and f, C and p are all defined over this subfield.

In the case f acts on the base as the identity, the proposition holds trivially. Assume that it is not that case. Let $O = L_\infty \cap F_\infty$. As in the proof of Proposition 6.2, we only have to consider the case $C \cap F_\infty = O$.

If a is a root of unity, we may replace f by f^n for some integer $n > 0$ and assume that $a = 1$ and $b = 1$. Since the transcendence degree of K is finite, we may embed K in the field of complex numbers $\mathbb{C}$. Let $|\cdot|$ be the standard absolute value on $\mathbb{C}$. Since f contracts F_∞ to O , there is a neighborhood U of O with respect to the usual euclidian topology such that for all point $q \in U \cap \{(x, y) \in \mathbb{C}^2 \mid \operatorname{Re}(x) > 0\}$, we have $\lim_{n \rightarrow \infty} f^n(q) = O$. Since $C \cap F_\infty = O$, there exists $M > 0$, such that $C \cap \{(x, y) \mid \operatorname{Re}(x) > M\} \subseteq U$ and we conclude by using Theorem 1.1 in this case.

If a is an algebraic number over $\mathbb{Q}$ and is not a root of unity, by [20, Theorem 3.8] there exists an absolute value $|\cdot|_v$ (either archimedean or non-archimedean) on $\mathbb{Q}$ such that $|a|_v > 1$. This shows that $(\mathbb{F}_m, f)$ satisfies the property DML by Proposition 6.2.

If a is not an algebraic number over $\mathbb{Q}$, we claim that there exists a field embedding $\iota : K \hookrightarrow \mathbb{C}$ such that $|\iota(a)| > 1$, and we may conclude again by using Proposition 6.2.

It thus remains to prove the claim. There is a subring R of K which is finitely generated over $\overline{\mathbb{Q}}$, such that f, C and p are all defined over R . There is an integer $l > 0$, such that $R = \overline{\mathbb{Q}}[t_1, \dots, t_l]/I$, where I is a prime ideal of $\overline{\mathbb{Q}}[t_1, \dots, t_l]$. It induces an embedding $\text{Spec } R := V \subseteq \mathbb{A}_{\overline{\mathbb{Q}}}^l$. We set

$$V_{\mathbb{C}} := V \times_{\text{Spec } \overline{\mathbb{Q}}} \text{Spec } \mathbb{C} \subseteq \mathbb{A}_{\mathbb{C}}^l.$$

For any polynomial $F \in \overline{\mathbb{Q}}[t_1, \dots, t_l] \setminus I$, we also define $V_F := \{F = 0\}$. Then $V_{\mathbb{C}} \setminus V_F$ is a dense open set in the usual euclidian topology. Since $\overline{\mathbb{Q}}[t_1, \dots, t_l] \setminus I$ is countable, the set $V_{\mathbb{C}} \setminus (\bigcup_{F \in \overline{\mathbb{Q}}[t_1, \dots, t_l] \setminus I} V_F)$ is dense. We view a as a nonconstant holomorphic function on $V_{\mathbb{C}}$, there is a open set $W \subseteq V_{\mathbb{C}}$ such that $|a| > 1$ on W .

Pick a closed point $(s_1, \dots, s_l) \in W \setminus (\bigcup_{F \in \overline{\mathbb{Q}}[t_1, \dots, t_l] \setminus I} V_F)$ and consider the unique morphism $\iota : R = \mathbb{Q}[t_1, \dots, t_l]/I \rightarrow \mathbb{C}$ sending t_i to s_i . This morphism is in fact an embedding. We may extend it to an embedding of K as required. $\square$

7. UPPER BOUND ON HEIGHTS WHEN $\lambda(f) > 1$

7.1. Introduction to heights. This section is a brief introduction to heights and their main properties. See [18] or [19] for detail. In this section, we denote by C a smooth curve defined over an algebraically closed field and by $K(C)$ its function field.

Proposition-Definition 7.1. Let K/L be a finite field extension where $L = \mathbb{Q}$ or $K(C)$, and $p \in \mathbb{P}^n(K)$ be a point with homogeneous coordinate $p = [x_0 : \dots : x_n]$ where $x_0, \dots, x_n \in K$. The *height* of p is the quantity

$$H_{\mathbb{P}^n}(p) := \left(\prod_{v \in \mathcal{M}_K} \max\{|x_0|_v, \dots, |x_n|_v\}^{n_v} \right)^{1/[K:L]}.$$

The height $H_{\mathbb{P}^n}(p)$ does not depend on the choice of homogeneous coordinates of p and the choice of field extension K which contains p .

When $L = K(C)$, we have a geometric interpretation of the height $H_{\mathbb{P}^n}(p)$. Observe that $\mathbb{P}_L^n$ is the generic fiber of the trivial fibration $\pi : \mathbb{P}_C^n := \mathbb{P}^n \times C \rightarrow C$. We set $s_p : D \rightarrow \mathbb{P}_C^n$ the normalization of the Zariski closure of p in $\mathbb{P}_C^n$. Then we have $H_{\mathbb{P}^n}(p) = e^{\deg(s_p^* \mathcal{O}_{\mathbb{P}_C^n}(1))}$.

Proposition 7.2. Let $k = \overline{\mathbb{Q}}$ or the algebraically closure of a function field of a curve. Let $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^m$ be a rational map and X be a subvariety of $\mathbb{P}_k^n$ such that $I(f) \cap X$ is empty and the restriction $f|_X$ is finite of degree d onto its image $f(X)$.

Then there exist $A > 0$ such that for all point $p \in X$, we have

$$\frac{1}{A} H_{\mathbb{P}^n}(p)^d \leq H_{\mathbb{P}^m}(f(p)) \leq A H_{\mathbb{P}^n}(p)^d.$$

Proposition 7.3 (Northcott Property). Let $K/\mathbb{Q}$ be a number field, and $B > 0$ be any constant. Then the set

$$\{p \in \mathbb{P}^n(K) \mid H_{\mathbb{P}^n}(p) \leq B\}$$

is finite.

7.2. Upper bound on heights. Let K be a number field or a function field of a smooth curve over an algebraically closed field. Let $f : \mathbb{A}_K^2 \rightarrow \mathbb{A}_K^2$ be any birational polynomial morphism defined over $\overline{K}$ and assume that $\lambda(f) > 1$.

According to Theorem 2.4, there exists a compactification X of $\mathbb{A}_K^2$, a point $Q \in X \setminus \mathbb{A}_K^2$ such that f extends to a birational transformation on X which satisfies the following properties up to a positive iteration.

- (i) The map f is algebraically stable on X .
- (ii) There exists a f -fixed point $Q \in X \setminus \mathbb{A}_K^2$, such that $df|_Q = 0$.
- (iii) We have $f(X \setminus \mathbb{A}_K^2) = Q$.

We fix an embedding $X \subseteq \mathbb{P}_K^N$. Let C be a curve in $\mathbb{A}_K^2$, we extend C to a curve in X .

Proposition 7.4. *Suppose that C is non periodic and $C \setminus \mathbb{A}_K^2 = \{Q\}$. Then there exists a number $B > 0$ such that for any point $p \in C(K)$ for which the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite, we have $H_{\mathbb{P}^N}(p) \leq B$.*

Proof. Assume that X, f, C and Q are all defined over K and $Q = [1 : 0 : \cdots : 0] \in \mathbb{P}_K^N$. There exists an element $a \in K^*$ and $F_i \in (x_1, \dots, x_N)K[x_1, \dots, x_N]$ for $i = 0, \dots, N$ such that

$$f([1 : x_1 : \cdots : x_N]) = [a + F_0 : F_1 : \cdots : F_N]$$

for any $[1 : x_1 : \cdots : x_N] \in X$. Since f is regular at Q and $a \neq 0$, there is a finite set $S \subseteq \mathcal{M}_K^0$ such that for any $v \in \mathcal{M}_K^0 \setminus S$, we have $|a|_v = 1$ and all coefficients of f are defined in O_v . Recall that we may endow X with a metric d_v , see Section 2.4.

For any $v \in \mathcal{M}_K^0 \setminus S$, set $r_v := 1$ and $U_v := \{x \in X(K) \mid d_v(x, Q) < 1\}$. Since $df|_Q = 0$, we see that for all $x \in U_v$, $d_v(f(x), Q) \leq d_v(x, Q)^2$, hence

$$\lim_{n \rightarrow \infty} f^n(x) = Q.$$

For any $v \in S$, set $r_v := |a|_v$ and $U_v := \{x \in X(K) \mid d_v(x, Q) < r_v\}$. We see that for all $x \in U_v$, $d_v(f(x), Q) \leq d_v(x, Q)^2/r_v$, and again it follows that

$$\lim_{n \rightarrow \infty} f^n(x) = Q.$$

For any $v \in \mathcal{M}_K^\infty$, since $df|_Q = 0$, there is $r_v > 0$ such that for any $x \in U_v := \{x \in X(K) \mid d_v(x, Q) < r_v\}$ we have $f(x) \subseteq U_v$ and

$$\lim_{n \rightarrow \infty} f^n(x) = Q.$$

If $p \in \bigcup_{v \in \mathcal{M}_K} U_v$, Theorem 1.1 shows that C is periodic and this contradicts our assumption. In other words, we have

$$p \in C(K) - \bigcup_{v \in \mathcal{M}_K} U_v.$$

Let

$$i : \tilde{C} \rightarrow C \subseteq X$$

be the normalization of C and pick a point $Q' \in i^{-1}(Q)$. There is a positive integer l such that lQ' is a very ample divisor of $\tilde{C}$. So there is an embedding $j : \tilde{C} \hookrightarrow \mathbb{P}^M$ for some $M > 0$ such that

$$Q = [1 : 0 : \cdots : 0] = H_\infty \cap \tilde{C}$$

where $H_\infty = \{x_M = 0\}$ is the hyperplane at infinity. Let

$$g : \tilde{C} \rightarrow \mathbb{P}^1$$

be a morphism sending $[x_0 : \cdots : x_M] \in \tilde{C}$ to $[x_0, x_M] \in \mathbb{P}^1$. It is well defined since $\{x_0 = 0\} \cap H_\infty \cap \tilde{C} = \emptyset$. Then g is finite and

$$g^{-1}([1 : 0]) = H_\infty \cap \tilde{C} = [1 : 0 : \cdots : 0].$$

By base change, we may assume that $\tilde{C}, i, j, g$ are all defined over K .

In the function field case, there is a smooth projective curve D such that $K = K(D)$; and in the number field case, we set $D = \text{Spec } O_K$.

We consider the irreducible scheme $\tilde{\mathcal{C}} \subseteq \mathbb{P}_D^M$ over D whose generic fiber is $\tilde{C}$ and the irreducible scheme $\mathcal{X} \subseteq \mathbb{P}_D^N$ over D whose generic fiber is X . Then i extends to a map $\iota : \tilde{\mathcal{C}} \dashrightarrow \mathcal{X}$ over D birationally to its image. For any $v \in \mathcal{M}_K^0$, let

$$\mathfrak{p}_v = \{x \in O_v \mid v(x) > 0\}$$

be a prime ideal in O_v . There is a finite set T consisting of those places $v \in \mathcal{M}_K^0$ such that ι is not regular along the special fibre $\overline{C}_{O_v/\mathfrak{p}_v}$ at $\mathfrak{p}_v \in D$ or $\overline{C}_{O_v/\mathfrak{p}_v} \cap H_{\infty, O_v/\mathfrak{p}_v} \neq \{[1 : 0 : \cdots : 0]\}$.

For any $v \in \mathcal{M}_K^0 - T \cup S$, let

$$\begin{aligned} V_v &= \{[1 : x_1 : \cdots : x_M] \in \tilde{C}(K) \mid |x_i|_v < 1, i = 1, \dots, M\} \\ &= \{[1 : x_1 : \cdots : x_M] \in \tilde{C}(K) \mid |x_M|_v < 1\}. \end{aligned}$$

We set $t_v := 1$ and $\Omega_v := \{[1 : x] \in \mathbb{P}^1(K) \mid |x|_v < 1\}$, then $V_v = g^{-1}(\Omega_v) \cap \tilde{C}(K)$.

For any $v \in T \cup S \cup \mathcal{M}_K^\infty$, by the continuity of i , there is $s_v > 0$ such that

$$i(V_v) \in U_v$$

where $V_v = \{[1 : x_1 : \cdots : x_M] \in \tilde{C}(K) \mid |x_i|_v < s_v, i = 1, \dots, M\}$. Since $g^{-1}([1 : 0]) = \{[1 : 0 : \cdots : 0]\}$, there exists $t_v > 0$, such that

$$g^{-1}(\Omega_v) \cap \tilde{C} \subseteq V_v$$

where $\Omega_v = \{[1 : x] \in \mathbb{P}^1(K) \mid |x|_v < t_v\}$.

Set $W := \{x \in \tilde{C}(\overline{\mathbb{Q}}) \mid i(x) \in X(K) \setminus \bigcup_{v \in \mathcal{M}_K} U_v\}$. Since i is the normalization, if $i(x)$ is a K -point of C and $x \notin I(i^{-1})$ then x is a K -point of $\tilde{C}$. So $j(x)$ is a K -point of $\mathbb{P}^M$ and $g(x)$ is a K -point of $\mathbb{P}^1$.

If $j(x) = [x_0 : \cdots : x_m] \in W \setminus i^{-1}(I(i^{-1}))$, then we have $x \in \tilde{C}(K) \setminus \bigcup_{v \in \mathcal{M}_K} V_v$, so that

$$g(x) \in \mathbb{P}^1(K) \setminus \left(\bigcup_{v \in \mathcal{M}_K} \Omega_v \right).$$

For any $y = [y_0 : y_1] \in \mathbb{P}^1(K) \setminus (\bigcup_{v \in \mathcal{M}_K} \Omega_v)$, we have $|y_1/y_0|_v \geq t_v$ for all v . Then we have

$$\begin{aligned} H_{\mathbb{P}^1}(y)^{[K:\mathbb{Q}]} &= \prod_{v \in \mathcal{M}_K} \max\{|y_0|_v, |y_1|_v\}^{n_v} \\ &\leq \prod_{v \in \mathcal{M}_K} \max\{|y_1|_v/t_v, |y_1|_v\}^{n_v} \\ &= \prod_{v \in \mathcal{M}_K} |y_1|_v^{n_v} \prod_{v \in \mathcal{M}_K} \max\{1, 1/t_v\}^{n_v} \\ &= \prod_{v \in \mathcal{M}_K} \max\{1, 1/t_v\}^{n_v} < \infty. \end{aligned}$$

Proposition 7.2 implies that there is a number $A > 0$, such that $H_{\mathbb{P}^M}(x) \leq A$ for all $x \in W \setminus i^{-1}(I(i^{-1}))$. We conclude by Proposition 7.2 that there is a number $B > 0$ such that $H_{\mathbb{P}^N}(p) \leq B$ for all points $p \in X(K) \setminus \bigcup_{v \in \mathcal{M}_K} U_v$. $\square$

8. PROOF OF THEOREM A

Let C be a curve in $\mathbb{A}_K^2$. We want to show that for any point $p \in \mathbb{A}^2(\overline{\mathbb{Q}})$ such that the set

$$\{n \in \mathbb{N} \mid f^n(p) \in C\}$$

is infinite, then p is preperiodic.

According to Section 6, we may assume that $\lambda(f) > 1$. As in Section 7.2, we use Theorem 2.4 to get a compactification X of $\mathbb{A}_K^2$. For simplicity, we write f for $\tilde{f}$ the map induced by f on X . There exists $n \geq 1$ such that f^n contracts $X \setminus \mathbb{A}_K^2$ to a supperattracting fixed point $Q \in X \setminus \mathbb{A}_K^2$. We extend C to a curve in X . Suppose that C is non periodic. By Theorem 1.2, we may assume that $C(K) \setminus \mathbb{A}^2(K) = \{Q\}$.

Fix an embedding $X \hookrightarrow \mathbb{P}_K^N$ for some $N \geq 1$.

We first treat the case $K = \overline{\mathbb{Q}}$.

There is a number field K such that both f and p are defined over K . Then $f^n(p) \in \mathbb{A}^2(K)$ for all $n \geq 0$.

Proposition 7.4 and the Northcott Property imply that the set $\{f^n(p) \mid n \geq 0\} \cap C$ is finite. Since the set $\{n \in \mathbb{N} \mid f^n(p) \in C\}$ is infinite, there exists $n_1 > n_2 > 0$ such that $f^{n_1}(p) = f^{n_2}(p)$. We conclude that p is preperiodic.

Next we consider the general case of an algebraically closed field K of characteristic 0.

Let $\mathbb{F}$ be the prime field of K . We may assume that $\text{tr.d.} K/\mathbb{F}$ of K over $\mathbb{F}$ is finite. We argue by induction on $\text{tr.d.} K/\mathbb{F}$.

If $\text{tr.d.} K/\mathbb{F} = 0$, then $K = \overline{\mathbb{Q}}$ and we have already treated.

If $\text{tr.d.} K/\mathbb{F} \geq 1$, then there is an algebraically closed subfield k of K such that $\text{tr.d.} k/\mathbb{F} = \text{tr.d.} K/\mathbb{F} - 1$.

There is a smooth projective curve D over k , such that X, f, p, Q and C are defined over the function field $K(D)$ of D .

We consider the irreducible scheme

$$\pi : \mathcal{X} \subseteq \mathbb{P}_D^N \rightarrow D$$

over D whose generic fiber is X and $\mathcal{C} \subseteq \mathbb{P}_D^N$ the Zariski closure of C in $\mathcal{X}$.

The map f extends to a birational map $f' : \mathcal{X} \dashrightarrow \mathcal{X}$ over D . For any $x \in D$, denote by X_x and C_x the fiber of $\mathcal{X}$ and of $\mathcal{C}$ at x respectively and denote by f_x the restriction of map f' to the fiber X_x .

Proposition 7.4 implies that there is a number $M \geq 0$ such that for all $n \geq 0$ either $f^n(p) \notin C$ or $H_{\mathbb{P}^N}(f^n(p)) \leq M$.

A point $s \in X_{K(D)}$ associates to its Zariski closure in $\mathcal{X}$ which is a section of $\pi : \mathcal{X} \rightarrow D$. For simplicity, we also write s for this section. Then the height of s is

$$H_{\mathbb{P}^N}(s) = e^{(s \cdot L)}$$

where $L := O_{\mathbb{P}_D^N}(1)$.

For any section s , observe that π induce an isomorphism from s to the curve D . We may consider the Hilbert polynomial

$$\chi(L|_s^{\otimes n}, s) = 1 - g(s) + n(s \cdot L) = 1 - g(D) + n \log H(s).$$

It follows that there is a quasi-projective k -variety M_H parameterizes the sections s of π such that $H_{\mathbb{P}^N}(s) \leq M$.

Let T_1 be the set of points $x \in D$ such that f_x is birational and $I(f_x^{-1}) \cap I(f_x) \neq \emptyset$. Observe that T_1 is finite. Let T_2 be the set of the points $x \in D - T_1$, such that C_x is fixed. Since C is not fixed, T_2 is finite. Because k is algebraically closed, $D - (T_1 \cup T_2)$ is infinite. For any point $x \in D$, denote by $p_x : M_H \rightarrow X_x$ the map sending s to $s(x)$. Pick a sequence of distinct points $\{x_i \in D - (T_1 \cup T_2)\}_{i \geq 0}$. For any $l \geq 1$, let

$$p_l = \prod_{i=1}^l p_{x_i} : M_H \rightarrow \prod_{i=1}^l X_{x_i}.$$

We see that any two points $s_1, s_2 \in M_H$ are equal if and only if $p_i(s_1) = p_i(s_2)$ for all $i \geq 0$.

We claim the following lemma, and prove it later.

Lemma 8.1. *Let X be any reduced quasi-projective variety over an algebraically closed field k . For any $i \geq 1$, let $\pi_i : X \rightarrow Y_i$ be a morphism. If for any different points $x_1, x_2 \in X$, there exists $i \geq 0$, such that $\pi_i(x_1) \neq \pi_i(x_2)$. Then for l large enough*

$$p_l = \prod_{i=1}^l \pi_i : X \rightarrow \prod_{i=1}^l Y_i$$

is finite.

By Lemma 8.1, there is an integer L large enough, such that the map p_L is finite. By Proposition 5.1, C_{x_i} is not periodic for all $i \geq 1$. The set $\{n \geq 0 | f^n(p) \in C\}$ is infinite, and we denote it by $\{n_i\}_{i \geq 0}$ and assume that $n_{i+1} > n_i$. For any $i \geq 0$, there exists $s_i \in M_H$ such that $s_i = f^{n_i}(p)$. By the induction hypothesis, we know

that $s_{n_0}(x_i) = f^{n_0}(p)(x_i)$ is a preperiodic point of f_{x_i} for any $1 \leq i \leq L$. Then the orbit O_i of $p(x_i)$ in X_{x_i} is finite. So the set

$$p_l(\{s_i\}_{i \geq 0}) \subseteq \prod_{i=0}^l O_i$$

is finite. Since p_l is finite, then we have $\{s_i\}_{i \geq 0}$ is finite. Then there is $i_1 > i_2$ such that $s_{i_1} = s_{i_2}$, and $f^{n_{i_1}}(p) = f^{n_{i_2}}(p)$. Then p is preperiodic. $\square$

Proof of Lemma 8.1. We prove this lemma by induction on the dimension of X .

If $\dim X = 0$, then the result is trivial.

If $\dim X > 0$, we may assume that X is irreducible. We pick any point $x \in X$, and let F_l be the fiber of p_l which contains x . Observe that

$$F_{l+1} \subseteq F_l,$$

so that there is an integer $L' \geq 1$, such that for any $L \geq L'$,

$$F_L = \bigcap_{l \geq 0} F_l.$$

Since for any point $x_1 \in X - \{x\}$, there exists $i \geq 0$, such that $\pi_i(x_1) \neq \pi_i(x)$, we have

$$F_L = \bigcap_{l \geq 0} F_l = \{x\},$$

so that

$$\dim X - \dim p_L(X) \leq \dim F_L = 0.$$

In particular p_L is generically finite. It means that there exists an open set U of $p_L(X)$, such that $p_L : p_L^{-1}(U) \rightarrow U$ is finite. Set $X' = X - p_L^{-1}(U)$, then we have $\dim X' \leq \dim X - 1$.

By the induction hypothesis, there is $L'' \geq L'$, such that for any $L \geq L''$, $p_L|_{X'}$ is finite and then p_L is finite. $\square$

REFERENCES

- [1] Shreeram Shankar Abhyankar. *Resolution of singularities of embedded algebraic surfaces*. Pure and Applied Mathematics, Vol. 24. Academic Press, New York, 1966.
- [2] Mauricio Velasco Antonio Laface. A survey on cox rings. arXiv:0810.3730.
- [3] J. P. Bell, D. Ghioca, and T. J. Tucker. The dynamical Mordell-Lang problem for étale maps. *Amer. J. Math.*, 132(6):1655–1675, 2010.
- [4] Jason P. Bell. A generalised Skolem-Mahler-Lech theorem for affine varieties. *J. London Math. Soc. (2)*, 73(2):367–379, 2006.
- [5] Robert L. Benedetto, Dragos Ghioca, Pär Kurlberg, and Thomas J. Tucker. A case of the dynamical Mordell-Lang conjecture. *Math. Ann.*, 352(1):1–26, 2012. With an appendix by Umberto Zannier.
- [6] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [7] Laurent Denis. Géométrie et suites récurrentes. *Bull. Soc. Math. France*, 122(1):13–27, 1994.
- [8] Jeffrey Diller and Charles Favre. Dynamics of bimeromorphic maps of surfaces. *Amer. J. Math.*, 123(6):1135–1169, 2001.

- [9] Tien-Cuong Dinh and Nessim Sibony. Une borne supérieure pour l'entropie topologique d'une application rationnelle. *Ann. of Math. (2)*, 161(3):1637–1644, 2005.
- [10] Gerd Faltings. The general case of S. Lang's conjecture. In *Barsotti Symposium in Algebraic Geometry (Abano Terme, 1991)*, volume 15 of *Perspect. Math.*, pages 175–182. Academic Press, San Diego, CA, 1994.
- [11] Charles Favre and Mattis Jonsson. Dynamical compactification of $\mathbf{C}^2$. *Ann. of Math. (2)*, 173(1):211–248, 2011.
- [12] S. Friedland and J. Milnor. Dynamical properties of plane polynomial automorphisms. *Ergodic Theory Dyn. Syst.*, 9:67–99, 1989.
- [13] William Fulton. *Algebraic curves. An introduction to algebraic geometry*. W. A. Benjamin, Inc., New York-Amsterdam, 1969. Notes written with the collaboration of Richard Weiss, Mathematics Lecture Notes Series.
- [14] D. Ghioca and T. J. Tucker. Periodic points, linearizing maps, and the dynamical Mordell-Lang problem. *J. Number Theory*, 129(6):1392–1403, 2009.
- [15] Dragos Ghioca, Thomas J. Tucker, and Michael E. Zieve. Intersections of polynomials orbits, and a dynamical Mordell-Lang conjecture. *Invent. Math.*, 171(2):463–483, 2008.
- [16] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [17] Ernst Kunz. *Introduction to plane algebraic curves*. Birkhäuser Boston Inc., Boston, MA, 2005. Translated from the 1991 German edition by Richard G. Belshoff.
- [18] J-P. Serre. *Lectures on the Mordell-Weil theorem*. Friedr. Vieweg & Sohn, Braunschweig, 1989. Translated from the French and edited by Martin Brown from notes by Michel Waldschmidt.
- [19] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1986.
- [20] Joseph H. Silverman. *The arithmetic of dynamical systems*, volume 241 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2007.
- [21] Paul Vojta. Integral points on subvarieties of semiabelian varieties. I. *Invent. Math.*, 126(1):133–181, 1996.

CENTRE DE MATHÉMATIQUES LAURENT SCHWARTZ ÉCOLE POLYTECHNIQUE, 91128, PALAISEAU
CEDEX, FRANCE

E-mail address: `jxie@clipper.ens.fr`